



VIT

Vellore Institute of Technology

REG.NO.:

SCHOOL OF ADVANCED SCIENCES
CONTINUOUS ASSESSMENT TEST - I
WINTER SEMESTER 2024-2025

SLOT: B1+TB1

Programme Name & Branch : B.Tech.
Course Code and Course Name : BMAT202L; Probability and Statistics
Faculty Name(s) : Common Slot QP
Class Number(s) : Common Slot QP
Date of Examination : 28-Jan-2025 (Session 1)
Exam Duration : 90 minutes

Maximum Marks: 50

General instruction(s):

- Answer All Questions
- M - Max mark; CO - Course Outcome; BL - Blooms Taxonomy Level (1 - Remember, 2 - Understand, 3 - Apply, 4 - Analyse, 5 - Evaluate, 6 - Create)
- Course Outcomes
CO1: Compute and interpret descriptive statistics using numerical and graphical techniques
CO2: Understand the basic concepts of random variables and find an appropriate distribution for analyzing data specific to an experiment
CO3: Apply statistical methods like correlation

Q. No	Question	M	CO	BL																				
1.	In a factory employing 3000 persons, 5 percent earn less than Rs. 3 per hour, 580 persons earn Rs. 3.01 to Rs. 4.50 per hour, 30 percent earn from Rs. 4.51 to Rs. 6.00 per hour, 500 persons earn from Rs. 6.01 to Rs. 7.50 per hour, 20 percent earn from Rs. 7.51 to Rs. 9.00 per hour, and the rest earn Rs. 9.01 or more per hour. What are the mean, median, and mode wage?	10	1	3																				
2.	The following table shows the distribution of employees based on their weekly working hours: <table border="1" style="margin-left: 20px;"> <tr> <td>Weekly hours</td> <td>30-34</td> <td>35-39</td> <td>40-44</td> <td>45-49</td> <td>50-54</td> <td>55-59</td> </tr> <tr> <td>No. of employees</td> <td>4</td> <td>6</td> <td>10</td> <td>14</td> <td>8</td> <td>8</td> </tr> </table> For the given data, calculate quartile deviation, standard deviation and compare the results to determine which measure of dispersion is better.	Weekly hours	30-34	35-39	40-44	45-49	50-54	55-59	No. of employees	4	6	10	14	8	8	10	1	2						
Weekly hours	30-34	35-39	40-44	45-49	50-54	55-59																		
No. of employees	4	6	10	14	8	8																		
3.	A random variable X has the following probability distribution: <table border="1" style="margin-left: 20px;"> <tr> <td>x</td> <td>-2</td> <td>-1</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>$p(x)$</td> <td>0.1</td> <td>k</td> <td>0.2</td> <td>$2k$</td> <td>0.3</td> <td>$3k$</td> </tr> </table> (a) Find the value of k , (b) Evaluate $P(X < 2)$ and $P(-2 < X < 2)$, (c) find the cdf of X and (d) evaluate the mean of X	x	-2	-1	0	1	2	3	$p(x)$	0.1	k	0.2	$2k$	0.3	$3k$	10	2	3						
x	-2	-1	0	1	2	3																		
$p(x)$	0.1	k	0.2	$2k$	0.3	$3k$																		
4.	The joint pdf of a two-dimensional RV (X, Y) is given by $f(x, y) = xy^2 + \frac{x^2}{8}$, $0 \leq x \leq 2, 0 \leq y \leq 1$ Compute (i) $P(X > 1)$, (ii) $P(Y < 1/2)$, (iii) $P(X > 1/Y < 1/2)$, (iv) $P(X < Y)$ and (v) $P(X + Y \leq 1)$.	10	2	4																				
5.	Find the coefficient of correlation between X and Y using the following data: <table border="1" style="margin-left: 20px;"> <tr> <td>X</td> <td>5</td> <td>10</td> <td>15</td> <td>20</td> <td>25</td> <td>30</td> <td>35</td> <td>40</td> <td>45</td> </tr> <tr> <td>Y</td> <td>16</td> <td>19</td> <td>23</td> <td>26</td> <td>30</td> <td>33</td> <td>36</td> <td>40</td> <td>43</td> </tr> </table>	X	5	10	15	20	25	30	35	40	45	Y	16	19	23	26	30	33	36	40	43	10	3	2
X	5	10	15	20	25	30	35	40	45															
Y	16	19	23	26	30	33	36	40	43															
