



VIT

Vellore Institute of Technology
(Approved to be a University under section 3 of the UGC Act, 1956)

Course: BMAT101L - Calculus

Class NBR(s): 5453 / 5456 / 6425

Time: Three Hours

Slot: D2+TD2

Max. Marks: 100

KEEPING MOBILE PHONE/SMART WATCH, EVEN IN 'OFF' POSITION, IS TREATED AS EXAM MALPRACTICE

Answer any TEN Questions

(10 X 10 = 100 Marks)

1. Determine the value of 'c' for which the function $f(x) = \sqrt{x(1-x)}$ in $[0,1]$ satisfies Mean Value theorem. Also check whether this function satisfies Rolle's theorem in $[0,2]$ [10]
2. Find the area of the region R bounded by the parabola $4y = x^2$ and the line $y = 2x$ in the first quadrant. If R is revolved about Y-axis to form a solid, find the volume of the solid. [10]
3. a) If $u = \tan^{-1}\left(\frac{y}{x}\right)$ where $x = e^t - e^{-t}$ and $y = e^t + e^{-t}$, find $\frac{du}{dt}$ [5+5]
b) Show that $u = 3x + 2y - z$, $v = x - 2y + z$ and $w = x(x + 2y - z)$ are functionally dependent.
4. Expand using Taylor series for $f(x, y) = \cos x \cos y$ at the origin upto 3rd order and hence find $f(0.1, 0.1)$. [10]
5. Use Lagrange multiplier method, to find the largest volume of the rectangular parallelepiped that can be inscribed in the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ [10]
6. Change the order of integration and evaluate $\int_0^1 \int_{x^2}^{2-x} xy \, dy \, dx$ [10]
7. Evaluate using spherical coordinates $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} \frac{dx dy dz}{\sqrt{1-x^2-y^2-z^2}}$ [10]
8. Using Beta and Gamma functions, evaluate a) $\int_0^{\frac{\pi}{2}} \sqrt{\cot \theta} \, d\theta$ and b) $\int_0^1 \frac{x dx}{\sqrt{1-x^5}}$ [5+5]
9. a) Determine the directional derivative of $\phi = 2xy - 3y^2$ at $(5,5)$ in the direction $4\hat{i} + 3\hat{j}$ [10]
b) Find $\text{div } \vec{F}$ and $\text{curl } \vec{F}$ for $\vec{F} = \text{grad}[x^3y + y^3z + z^3x - x^2y^2z^2]$
10. Show that $\vec{F} = (2x + yz)\hat{i} + (4y + zx)\hat{j} - (6z - xy)\hat{k}$ is both solenoidal and irrotational. Find also its scalar potential. [10]
11. Verify Green's theorem for $\oint_C [3x^2 - 8y^2]dx + [4y - 6xy]dy$ where C is the boundary of the region $x = 0, y = 0$ and $x + y = 1$ [10]
12. Using Stoke's theorem evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F} = y\hat{i} + z\hat{j} + x\hat{k}$ and C is the boundary of the plane $3x + 2y + z = 6$ in the first quadrant. [10]

grad n
grad

Handwritten calculations and diagrams for question 12, including a diagram of a triangle in the first quadrant and various algebraic steps.